



NUTRITIONAL SOLUTIONS THAT HELP YOU MOVE FORWARD

ALEMA KOUDIJS FEED PLC (AKF) IS A PROFESSIONAL ANIMAL FEED PRODUCTION COMPANY AND THE MOST MODERN IN ETHIOPIA. AKF IS A JOINT VENTURE WITH DE HEUS ANIMAL NUTRITION.

WHAT MAKES OUR JOINT VENTURE UNIQUE?

1. Using 100 years experience from our Dutch partner De Heus Animal Nutrition
2. Quality assured through laboratory analysis
3. On-the-farm support and advise
4. Improving farmers profit by increasing production

AKF offers you the right feed for each stage of your broiler chicken. Each type of feed is perfectly balanced to meet the demand of the broiler during the specific period.

CRUMBLE AND PELLET FOR UNBEATABLE RESULTS!

All our broiler feeds are pelletized, which is unique for Ethiopia. Why pelletizing:

- Higher feed intake and thus faster growth; more cycles per year
- More efficient growth; less kg of feed per kg of meat
- No feed waste
- No selective eating

CHICKEN STARTER CRUMBLE (21% CRUDE PROTEIN, 3050 KCAL)

Chicken Starter Crumble is the optimum feed to give your chicks a good starting growth. The growth in the starter period is decisive for later growth. During this stage, the chick is developing its organs and thus need a lot of protein and energy. The 7 day weight is very important: Every 1 gram higher body weight at 7 days, will result in 5 grams extra body weight at slaughter age.

BROILER GROWER CRUMBLE (19% CRUDE PROTEIN, 3150 KCAL)

After 10 days, we change to Broiler Grower Crumble. This feed is slightly lower in protein and higher in energy.

BROILER FINISHER FEED (18% CRUDE PROTEIN, 3250 KCAL)

After 30 days, we change to Broiler Finisher Feed. This feed is slightly lower in protein and higher in energy.

All feeds include coccidiostatics, there is no need of giving additional.

AGE IN DAYS	TYPE OF FEED	FEED INTAKE DURING PERIOD
0-10 days	Chicken Starter Crumble	250 gr
11-30 days	Broiler Grower Crumble	2 kg
30 days-slaughter	Broiler Finisher Feed	2 kg

DAY	ROOM TEMPERATURE °C
0	33
3	32
7	30
14	27
21	24
28	22
35	20
42	19

BASIC GUIDELINES

- Feed minimum 3 times per day, never limit the feed intake; More feed intake = More growth
- Fresh water should always be available
- Follow the temperature closely and make sure there is sufficient ventilation

CONTROL CHART BROILERS



FEED TYPE	DAY	BODY WEIGHT (G) TARGET	BODY WEIGHT ACTUAL (G)	CUM. INTAKE TARGET	CUM. INTAKE ACTUAL	FCR	MEDICATION
STARTER	0	42					
	1	48		11			
	2	58		13			
	3	72		15			
	4	88		18			
	5	106		22			
	6	126		25			
	7	147		30			
	8	172		34			
	9	199		38			
	10	228		42			
GROWER	11	259		46			
	12	293		51			
	13	330		55			
	14	369		59			
	15	410		66			
	16	454		70			
	17	501		75			
	18	549		80			
	19	600		86			
	20	653		91			
	21	708		96			
	22	765		103			
	23	824		107			
	24	885		113			
	25	948		118			
	26	1012		122			
	27	1076		129			
	28	1144		133			
	29	1212		139			
	30	1281		143			
FINISHER	31	1350		148			
	32	1421		153			
	33	1493		157			
	34	1565		161			
	35	1637		165			
	36	1710		169			
	37	1783		173			
	38	1856		176			
	39	1929		180			
	40	2002		183			
	41	2075		186			
	42	2148		189			
	43						
	44						
	45						
	46						